



Brief Alcohol and Drugs Screening Tools 2 (B.A.D.S.T. 2)



screening tools





















Brief Alcohol & Drugs Screening Tool (BADST-2)



Each part of the BADST-2 is in itself a brief intervention, you are not completing this tool because it is a requirement, the tool is to be used when something has given you cause for concern in one of the three areas. (Alcohol, Drugs or Gambling). The tool is provided to enable you to structure a discussion around a difficult subject, find out more, and provide appropriate levels of reinforcement, advice or support, including brief interventions or referral to medical providers or specialist services. You will need the person's consent for each individual set of questions, you will also need consent to share the results as part of any referrals that follow.

Contents

page 1: BADST-2 combines three short screening tools on one page, these are:

AUDIT-C 3 questions that screen for alcohol related issues.
 DUDIT 3 questions which screen for drug use related issues.
 PGSI mini-screen 3 questions that screen for gambling related issues.

page 2: Instructions on how to score each section and what to do according to the score.

- If the brief screening on page 1 does not make you aware of any issues, then you do not need to go any further – that's your screening done
- If it does highlight any issues, then follow the guidance on page 2
- The following pages provide more detailed screening and assessment tools to use if suggested by a score on page one
- **AUDIT-C** short version, a quick reference guide to units of alcohol, scoring guide and instructions on actions to take.
- page 5: The Drug Use Screening Tool (DUST), identifies substances used, twelve screening questions, scoring guide and instructions on actions to take.
- **Problem Gambling Severity Index (PGSI)** screening tools, scoring guide and instructions on actions to take.
- page 11: Tools for Change, Building Motivation and Setting SMART Goals

Appendix 1: DECCA referral form

Printing Guide

The most eco-friendly way to use this tool is on a touch screen device with a stylus pen, however the most user-friendly way will be with a pen on paper; so when printing paper versions to use, please consider what you are concerned about and **only print pages that you may use**.

- In each section of this tool, only certain pages need to be printed, the other pages are for instruction only. Therefore:
 - p1 CRAST
 - p3 AUDIT
 - p6-8 DUST
 - p10 PGSI
- are the only pages you may need to print in the tool, they do not need to be printed in colour,
 please print in B&W and only print what you need.

The referral form for DECCA is included as **appendix 1**, only print if needed.

Brief Alcohol & Drugs Screening Tool (BADST-2)



Name:					
AUDIT: Alcoh	nol use in the pas	t 6 months			
1. How often do	o you have a drink	containing a	lcohol?		
Never □0	Monthly or less □1	Weekly □2	2-3 times a week □3	Daily □4	
2. How many u	nits of alcohol do	you drink typi	cally when you	are drinking?	
1-2 □0	3-4 □1	5-6 □2	7-9 □3	10+ □4	
3. Have you had Never □0	d 6 or more units Occasionally □1		more (m), on a s Weekly □3	single occasion? Daily □4	
			AUD	IT Score:	\supset
1. How often ar	lem drug use in the re you heavily und casionally □1	-		Daily □4	0
	ave you felt that yeasionally □1		ugs to feel well Weekly □3	or better? Daily □4	\bigcirc
•	t done something asionally □1	-	ave done becau Weekly □3	ıse of using druç Daily □4	gs?
			DUD	IT Score:	
PGSI: Gam	bling in the last 1	2 months			
never	= zero; sometimes	= one; most o	of the time = two	; almost always	= three
1. Have you bet	more than you coul	ld really afford	to lose?		
	criticized your bettin of whether or not yo	•	,	nbling problem,	Ŏ
3. Have you felt	guilty about the wa	y you gamble d	or what happens	when you gamble	∍?
			PG	SI Score:	<u> </u>

Brief Alcohol & Drugs Screening Tool (BADST-2)



The BADST-2 is three screening tool for alcohol use, problem drug use and gambling in the past six to twelve months.

AUDIT: Alcohol Use in the past 6 months

The Alcohol Use Disorders Identification Tool (AUDIT) was developed by the World Health Organisation (WHO) and has been used in a variety of health and social care settings.

A score of <u>less than 5</u> indicates lower risk drinking, an opportunity to raise awareness and give positive reinforcement. <u>Scores of 5+</u> require further assessment with full AUDIT for Alcohol (p3). This is also available as an online test <u>WHO/ Europe | Alcohol use - Take the AUDIT test now</u>

DUDIT: Problem drug use in the past 6 months

The 3 screening questions are adapted from The Drug Use Disorders Identification Test (DUDIT) manual. Bergman, A.H., et al. 2003,

A score of <u>less than 5</u> indicates lower risk drug use, an opportunity to give brief harm reduction advice and positive reinforcement. <u>Scores of 5+</u> require further assessment with the full DUST for substance use (p9).

PGSI: Gambling in the last 12 months

- Non-problem gambler Gamblers who gamble with no negative consequences
- 1 Low-risk gambler Gamblers who experience a low level of problems with few or no identified negative consequences
- 2-3 Moderate-risk Gamblers who experience a moderate level of problems leading to some negative consequences
- 4+ Problem gambler Gambling with negative consequences and a possible loss of control

Download information leaflets at: https://www.ourguideto.co.uk/decca-big-guide-to

DECCA Awareness Club: www.ourguideto.co.uk/decca-awareness-club/



Alcohol Use Disorders Screening Tool (AUDIT)



Name:	W orker:	 Date: /	/	,
	** • • • • • • • • • • • • • • • • • •	 /		

The Alcohol Use Disorders Identification Tool (AUDIT) was developed by the World Health Organisation (WHO) and modified for use in the UK and has been used in a variety of health and social care settings. Alcohol use screening tests - GOV.UK (www.gov.uk)

AUDIT-C Questions			Scoring			Your
	0	1	2	3		score
How often do you have a drink containing alcohol?	Never	Monthly or less	2-4 times a month	2-3 times a week	4+ times a week	
How many units of alcohol do you drink on a typical day when you are drinking?	1-2	3-4	5-6	7-9	10+	
How often have you had 6 or more units if female, or 8 or more if male, on a single occasion in the last year?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
				Tot	al score:	

A score of <u>less than 5</u> indicates lower risk drinking (see overleaf). <u>Scores of 5+</u> require the following 7 questions to be completed: For AUDIT translations in 40 languages: AUDIT translations (auditscreen.org)

AUDIT Questions			Scoring			Your
	0		2	3		score
How often during the last year have you found that you were not able to stop drinking once you had started?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
How often during the last year have you failed to do what was normally expected from you because of your drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
How often during the last year have you needed an alcoholic drink in the morning to get yourself going after a heavy drinking session?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
How often during the last year have you had a feeling of guilt or remorse after drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
How often during the last year have you been unable to remember what happened the night before because you had been drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily	
Have you or somebody else been injured as a result of your drinking?	No		Yes, but not in the last year		Yes, in the last year	
Has a relative or friend, doctor or other health worker been concerned about your drinking or suggested that you cut down?	No		Yes, but not in the last year		Yes, in the last year	
				Tot	al score:	

AUDIT	_	- AUDIT	_		0.00.0.
SCORING: ADD th	ne 2 scores	toaether to	o identify	necessarv	action



Alcohol Use Disorders Screening Tool (AUDIT)



https://alcoholchange.org.uk/alcohol-facts/interactive-tools/check-your-drinking/alcohol-units

AUDIT score intervention guide

AUDIT Score	Risk Category	Desired Action
0-4	Lower Risk	Positive reinforcement of low risk drinking guidelines
5-9	Increasing Risk	Brief intervention, reinforce low risk guidelines and explore strategies for cutting down
10-14	Higher Risk	Extended Brief Intervention and / or referring to local services for Brief Treatment.
15+	High Risk and Possible Dependence	Refer to specialist treatment services, if refused give safer drinking tips & use brief motivational interventions to promote treatment.

- **0-4 Simple Brief Advice:** An opportunity to educate people about low risk drinking levels and the risks of excessive alcohol use. **NB:** It is never safe to drink alcohol at all during pregnancy.
- **5-9 Brief Intervention to Reduce Use:** Person-centered discussion that uses motivational enhancement concepts to raise an individual's awareness of their substance use and enhance their motivation to change behaviour. Brief interventions are typically 5-15 minutes and should occur in the same session as the initial screening. Repeated sessions are more effective than a one-time intervention. The recommended behaviour change is to cut back to low risk drinking levels unless there are other medical reasons to abstain (liver damage, pregnancy, medication, etc.).
- **10-14 Brief Intervention to Reduce or Abstain (Brief Treatment if available) & Follow-up:** People with numerous or serious negative consequences from their alcohol use, or people who likely have an alcohol use disorder who cannot or are not interested in obtaining specialised treatment, should receive more numerous and intensive Brief Interventions with follow up. The recommended behaviour change is to cut back to low risk drinking levels or abstain from use. Brief treatment is 1 to 5 sessions, each 15-60 minutes. Refer for brief treatment if available (see below), if brief treatment is not available, secure follow-up in 2-4 weeks.
- **15+ Brief Intervention to Accept Referral:** The focus of the brief intervention is to enhance motivation for the person to accept a referral into treatment. If accepted, the provider should use a proactive process to facilitate access to specialty substance use disorder treatment for assessment and, if warranted, treatment. The aim is for the person to reduce use and accept the referral.

Refer to DECCA in Sandwell (appendix 1) who offer brief treatment, groupwork, and one-to-one support to reduce drinking, as well as interventions to manage health problems caused by drinking.

There are now free apps that some people may find useful for monitoring their alcohol consumption such as: <u>Drink Free Days</u> and <u>Dry January</u>



ALTHOUGH many young people will try drugs and alcohol at some time, most do not progress beyond experimentation. However, research indicates that many factors can increase the risk of a young person moving from 'drug & alcohol use' to 'drug & alcohol misuse', whilst some protective factors can reduce these risks. Unless you are a specialist drug & alcohol worker it can be difficult to distinguish between use and misuse, and to accurately assess these risk factors. This tool should help.

To complete this form, you do not need a comprehensive knowledge of drugs & alcohol but you may need to know how to contact your nearest drugs & alcohol service for young people (see opposite page). This service will be able to provide appropriate information, leaflets and guidance.

DUST is designed for use with young people about whom there may be concerns regarding drug / alcohol use.

- It is not a comprehensive drug/alcohol use assessment.
- It will indicate when specialist advice should be sought.
- It will help identify risk factors.

Defining the terms

Drug/alcohol use The consumption of a drug or alcohol by a young person. When the term 'use' is contrasted with 'misuse', 'use' means the consumption of a drug or alcohol that does not cause any perceptible immediate harm - even though it may carry some risk of harm.

Drug/ alcohol misuse Use of a drug / alcohol or combination of drugs and alcohol, that harms health or social functioning - either dependent use physical or psychological) or use that is part of a wider spectrum of problematic or harmful behaviour.

Vulnerable group Young people are at increased risk of drug / alcohol misuse if they belong to certain groups and this risk increases if there is membership of more than one group.

Protective factors Increase a young person's resilience to the development of drug / alcohol misuse problems.

Risk factors Increase the likelihood that drug / alcohol misuse will occur.

Protective Factors

- Positive temperament
- Intellectual ability
- Supportive family environment
- Social support system
- Caring relationship with at least one adult
- In education / employment / training

Belonging to a 'vulnerable' group

- Young sex workers
- Young offenders
- Looked after children
- Mental health problems
- School non-attenders
- Drug / alcohol misuse by parents
- Abuse within the family
- Homeless

Risk Factors

Social & Cultural Factors

- High levels of neighbourhood crime
- High levels of poverty & decay
- Easy drug availability
- Areas where there is widespread social acceptance of drug use
- Lack of perception of the risks from drugs / alcohol

Interpersonal & Individual Risk Factors

- Physiological & psychological factors
- Family dysfunction
- Behavioural difficulties
- Academic problems
- Association with peers who use drugs / alcohol
- Early onset of drug or alcohol use.

This form is a screening tool to indicate whether an assessment by a specialist service is needed. The Young Persons' Drug and Alcohol Service will use this information to determine an appropriate course of action. This may result in the young person being offered some form of intervention/treatment or in further advice and guidance being offered to the referrer. All information on completed DUST forms sent to DECCA will be treated in accordance with their confidentiality policies - a copy of which is available on request. (Exceptions to confidentiality include following Safeguarding procedures).

This tool is designed for two main purposes:

- To help people who have to make decisions about how to respond to drug / alcohol use by a young person.
- To allow services to create a caseload profile and audit the prevalence of drug / alcohol use within their case load.

The form is divided into sections designed to assess risk factors regarding:

- Drug and alcohol use
- Social situation/ behaviour
- General and psychological health

Instructions

- Complete the form by ticking the most appropriate responses. If in doubt, do NOT tick.
- A scoring system is employed for each section. The scores should be added up and the total written below each section. Please do not include past use in your scores.
- Once you have completed each section, refer to the scoring table.
- If the young person does not meet the threshold for referral but you still have concerns, please contact DECCA for advice and information.



DUST	Se		1:	
Drug	and	Alco	hol	Use

DUST Section 1:	,	,	
Drug and Alcohol Use	ن کی ا	USED	PAS
Drug Type	SCORE	USED IN LAST	PAST USE
Alcohol (booze)	2		
Amphetamine (speed, base, whiz, billy)	3		
Cannabis (marijuana, weed, skunk, dabs, shatter, oil)	2		
Cocaine (coke, charlie)	4		
Crack (rock, white)	7		
Ecstasy / MDMA / 2CB (pills, mandy)	4		
GHB	4		
Heroin (smack, scag, gear)	7		
IPED's (image and performance enhancing drugs) steroids, slimming tablets,	2		
tanning drugs			
Ketamine (ket, K, special-k)	4		
LSD (acid, trips, tabs)	4		
Magic Mushrooms (shrooms, mushies, caps)	4		
Nitrous Oxide (laughing gas)	2		
Novel Psycho-active Substances, NSP's (spice, M-cat, mamba)	4		
Psychoactive Compounds (salvia, DMT, ayahuasca)	3		
Solvents / Gas / Aerosols (glue, thinners, nail polish)	7		
Tobacco (cigarettes, baccie, fags)	1		
Other(s) please list (include misuse of over the counter or prescribed drugs). Score	2 2		
each:			
Drug/ Alcohol Use - Frequency			
Occasionally using drugs / alcohol	1		
Regular drug/alcohol use	2		
Injecting			
Not injecting	0		
Currently / recently injecting	5		
Drug/ Alcohol Use - Intoxication			
Drug / Alcohol use without loss of consciousness or aggression	0		
Drug / Alcohol use with loss of consciousness or aggression	5		
		1	
Contact with Other Drug/Alcohol users			
No drug / alcohol using friends Some friends who use drugs / alcohol and some who don't	1		
All friends use drugs / alcohol	2		
/ III THO TOS USE CITUYS / ALCOHOL			

10 mg (10 mg)	The second second	Control of the Control
		ı ı score:

Section 1	0-4	5-6	7+
Drug / Alcohol	Give brief drug	Refer to DECCA Young People's	Refer to DECCA Young People's
Use	and alcohol	Substance Misuse Service for more	Substance Misuse Service for
	information &	detailed drug and alcohol information &	further assessment and specialist
	advice (IBA)	advice (EIBA)	drug and alcohol support.

Drug Use Screening Tool (DUST)



DUST Section 2:

Social Situation / Behaviour

Family Drug / Alcohol Use		
No known family drug / alcohol misuse	0	
Known parental or close family member drug / alcohol misuse carer(s).	2	
Significantly affected by someone else's drug / alcohol misuse	5	
Living Situation		
No problems with accommodation	0	
Problems with accommodation, insecure or inadequate housing	1	
Looked after by Local Authority	2	
Homeless	6	
Adult Support		
Has supportive relationships with more than one adult	0	
Has supportive relationship with one adult	1	
Has no supportive relationship with adults	2	
Problematic Relationships (e.g. domestic violence at home)	2	
Education, Employment and Training		
In education / employment / training	0	
Truanting from school / risk of school exclusion / drug/alcohol related absences	1	
from school or work		
School excluded / unemployed	2	
Criminal Involvement		
No criminal involvement	0	
At risk of involvement in the Criminal Justice System	1	
Involved in Criminal Justice System or committing more serious crimes	2	
Sexual Behaviour		
Age appropriate / safe sexual behaviour	0	
Inappropriate behaviour / unsafe sexual behaviour	2	
Child Sexual Exploitation / child abuse/ abusive sexual relations	6	
Financial Name to the cloth		
No problems with debt Drug related debt	6	
Drug related debt	U	

Total section 2 score:

Section2	0-1 - Low	2-5 Medium	6+ High
Social	Low risk	Provide brief drug and alcohol	Refer to DECCA for more detailed
Situation /		information & advice and seek	drug and alcohol information &
Behaviour		further guidance / support from	advice. Consider a referral to
		DECCA.	Sandwell Children's Trust.
Continua			
Section3	0-4 Low	5-9 Medium	10+ High
General &	0-4 Low Low risk	Frovide brief drug and alcohol	10+ High Refer to DECCA for more detailed
	0 1 2011		
General &	0 1 2011	Provide brief drug and alcohol	Refer to DECCA for more detailed



DUST Section 3:

General Health				
No significant health problems repo	rted	0		
Teeth problems				
Stomach problems				
Regular headaches		1		
Difficulty sleeping		1		
Chronic fatigue		5		
Severe sleep problems		5		
Self-neglect		5		
Extreme weight loss		10		
Blackouts and / or memory loss		10		
Pregnant		10		
Fitting		10		
Accidental / planned overdose		10		
Psychological Health				
No significant psychological problem	ns	0		
Low self-esteem		1		
Mild anxiety		11		
Shyness		1		
Eating disorder / marked change in eating pattern (e.g. loss of appetite / bingeing)				
Frequent bouts of unhappiness / depression				
Self-harm				
Severe anxiety / panic attacks				
Suicide attempts				
Severe paranoia				
Hallucinations (when not under the influence of substances)				
Transconations (when not under the	initiative of substances)	10		
	Total section 3 s	score:		
Young Person's Details				
Name:	Age:			
Mobile #:	Sex:			
	- Com			
W orker's Details				
Name:	Position:			
Mobile #:	Organisation:			
Nort Change	, ~ 1			
Next Steps:	V.D. cianahara			
□ No Action	Y.P. signature:			
□ Repeat & Review in months□ Brief Intervention & Awareness□ Date://				
 □ Referral to DECCA 	/	• • • •		
☐ Safeguarding	W orker's signature:			

The DUST was developed for the Kent and Medway Drug Action Team by Neil Hunt, 2002. It is in the public domain.



The short-form Problem Gambling Severity Index

ivan	ne: Date://				
This instrument is formed of three questions from the PGSI, which are scored on a 4-point scale from never to almost always. It is asked to all participants of a survey who have gambled at least once in the last 12 months.					
P	PGSI: In the last 12 months				
	never = zero; sometimes = one; most of the time = two; almost always = three				
1	Have you bet more than you could really afford to lose?				
2	2. Have people criticized your betting or told you that you had a gambling problem, regardless of whether or not you thought it was true?				
3	3. Have you felt guilty about the way you gamble or what happens when you gamble?				
	PGSI Score:				

Scoring instructions

- Non-problem gambler Gamblers who gamble with no negative consequences.
- 1 Low-risk gambler Gamblers who experience a low level of problems with few or no identified negative consequences.
- 2-3 Moderate-risk Gamblers who experience a moderate level of problems leading to some negative consequences.
- 4+ Problem gambler Gambling with negative consequences and a possible loss of control.

The short-form Problem Gambling Severity Index (PGSI mini-screen) was developed for the Commission from the full 9-item PGSI by Dr. Rachel Volberg, 2012.

https://www.gamblingcommission.gov.uk/statistics-and-research/publication/problem-gambling-screens

Online Self-assessment tool

Self-assessment tool - GamCare

If you are not sure how much gambling has become a problem for you, you can take an online assessment to find out how much of an impact gambling is having in your life.

The assessment will give you a series of statements and ask you to select how much the statement applies to your gambling behaviour on a scale of 1 - 10.

At the end of the test you will be presented with a breakdown of how gambling is affecting your life and will give you personalised recommendations for your next steps.



Problem Gambling Screening Inventory

Name:	_/	_/ _	
-------	----	------	--

Used in the Health Survey for England, Scottish Health Survey, and the Welsh Problem gambling Survey. The PGSI consists of nine items and each item is assessed on a four-point scale: never, sometimes, most of the time, almost always. Responses to each item are given the following scores:

Thinking about the last twelve months never = 0; sometimes = 1; most of the time = 2; almost always = 3	Score
Have you bet more than you could really afford to lose?	
Have you needed to gamble with larger amounts of money to get the same feeling of excitement?	
When you gambled, did you go back another day to try to win back the money you lost?	
Have you borrowed money or sold anything to get money to gamble?	
Have you felt that you might have a problem with gambling?	
Has gambling caused you any health problems, including stress or anxiety?	
Have people criticized your betting or told you that you had a gambling problem, regardless of whether or not you thought it was true?	
Has your gambling caused any financial problems for you or your household?	
Have you felt guilty about the way you gamble or what happens when you gamble?	
Total:	

Scoring instructions

0	Non-problem gambler - Gamblers who gamble with no negative consequences.					
1-2	Low-risk gambler - Gamblers who experience a low level of problems with few or					
	no identified negative consequences.					
3-7	Moderate-risk - Gamblers who experience a moderate level of problems leading					
	to some negative consequences. Give encouragement to explore online resources					
	and a brief intervention to reduce or abstain from gambling.					
8+	Problem gambler - Gambling with negative consequences and a possible loss of					
	control. Brief intervention to reduce and accept referral to a specialist service.					

https://www.gamblingcommission.gov.uk/statistics-and-research/publication/problem-gambling-screens

When scores to each item are summed, a total score ranging from 0 to 27 is possible. A PGSI score of eight or more represents a problem gambler. This is the threshold determined by the developers of the PGSI.

Help with gambling problems

In the West Midlands the organisation commissioned by GamCare is Aquarius who provide support for people 16 years and over affected by gambling, either their own or that of a family member or friend, through 1-2-1 or group support sessions.

People can self-refer to this service or be referred by a professional. To find out more or refer call: 0300 456 4293 or email: gambling@aquarius.org.uk




W hat do you like about this behaviour, what do you get from it?	
W hat are your thoughts about this behaviour in the past, the present, and the future?	


In these boxes, weigh up the pros and cons of keeping things as they are, or trying to change them.

Staying a Advantages	s things are Disadvantages
	•
	•
	•
	•
	•
	•
	•
	•
	•
	•
	•

Making a change								
Advantages Disadvantages								
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							
	•							

List your three main reasons for changing this behaviour in the boxes below, then write down why each reason is important to you.










By When? Specific Actions - 3 small SMART steps 1. 2. 3. Strengths I have or need Helpful people & useful thoughts My Goal How will I know this has happened, what will be different? Possible problems and solutions Preparing for an Appointment Who is it with? Where is it exactly? When is it, at what time?



Appendix 1: DECCA Referral Form



Please complete this form in full, if this does not occur we may need to return the referral so further information can be added. The information supplied is for use of the DECCA Team and its contents are kept confidential. This document is retained at DECCA.

Email referrals as a WORD DOC to:

Decca_Team@sandwellchildrenstrust.org or call 0121 569 2201 / 07500 785889

Young Persons	Details:					
First Name(s)		Surname				
Date of Birth		Age				
Ethnicity		Sex	M/F			
Address:						
Contact number(s) Optional						
How does the y	young person want to be contacted?	•	Any disabilities, special educational needs or special			
Where does the	e young person want to be seen?		requirements?			
What school/er	mployment do they attend?					
GP Details (essential when making a referral)						
Is the Young P	erson involved with early help servi	ces or child	ren's social care? Yes / No			
If YES, contact	the worker to make them aware of t	his referral.				
Contact details for worker:						
Details of the early help intervention or what plan they are on:						
In order to support them, DECCA will need to keep details of any young person referred to their service, on their secure electronic system. You will need to seek their consent to the information on this form, being shared with DECCA. They will need to understand that DECCA will hold the information securely and retain the form for DECCA use only. They also need to understand that they can withdraw their consent to this referral at any time. More information on this can be found at https://www.sandwellchildrenstrust.org/privacy/ . Please ask the young person you are referring, to sign this form to agree to this, or if they have consented to the referral verbally, please add your signature so you can confirm on their behalf. For more information on how long records are kept, please speak to a DECCA Worker.						
Signed:		Date:				





Substance(s)/drug(s) used:
Please detail why you are referring this young person to DECCA and what you hope that they
gain from working with DECCA?
What does the young person want to achieve by working with DECCA?
what does the young person want to achieve by working with blook:

Appendix 1: DECCA Referral Form



Referrers Details:

What to tell young people	First Name			Surname		
about DECCA: We are	Job Title if applicable			Relationsh young pers	•	
confidential	Address					
We can arrange to meet you in a						
place that suits you best	Email:					
We won't try	Contact Numl	ber(s)				
and tell you what to do	Lead Professi this young pe					
We know quite a bit about drink and drugs	including <u>con</u> number and e not yourself					
We know	Is the young p	oerson's	parent(s) or carer	<u>(s)</u> aware of	this referral?	YES/NO
about what young people	Who has pare	_	oonsibility?			
in Sandwell do because	Contact detai	ls:				
we talk to them						
everyday	Note: If the refe	erral is to	be used for part of t	he exit strate	eav. for the you	ng person.
If your problem isn't drink or drugs	Note: If the referral is to be used for part of the exit strategy, for the young person, then the referral should be received in no less than 3 weeks before their intervention with you ends. If this does not occur then the referral may not be taken.					
we can help you get the	Ever admitted	l to hosp	ital for substance/	alcohol mis	suse?	YES/NO
right help If YES, give details of admission including: date/ who present/ CAMHS We will work					esent/ CAMHS r	eferral etc.
with you for as long as is needed but no						
longer than you want – we want to	Referrers sign	nature:				
provide the right service for YOU	Date referral o	complete	d:			
	Date referral i	eceived	by DECCA (DECC	A use only)	:	





Whether or not you are making a referral to DECCA, to evaluate this collection of practice tools, we would appreciate your feedback and welcome suggestions for future versions.

1.	Which parts of the BADST-2 have you used?					
□ BADST Mini Screens (p1)□ PGSI (p9)□ AUDIT (p3)□ Building Motivation (p1						
					(p11)	
	□ DUST - section 1 (p6)	Setti	ng Goa	ls (p12)	
	□ DUST - section 2 (p7)		CA Ref	erral Fo	rm (p13	3)
	□ DUST - section 3 (p8)					
2.	How have you used this tool?					
	☐ In a practice situation with a young person					
	☐ With a colleague in peer reflection					
	☐ By reflecting on how to use it in your practic	e				
3.	Do you agree or disagree with these statemer	nts?				
		Disagree				Agree
	The instructions were easy to follow.	1	2	3	4	5
	The tool was easy to use.	1	2	3	4	5
	The tool maintained focus on the topic.	1	2	3	4	5
	The tool aided communication.	1	2	3	4	5
	The tool opened up a wider conversation.	1	2	3	4	5
	The tool supported awareness raising.	1	2	3	4	5
	The tool helped in planning the next step.	1	2	3	4	5
4.	Is there any part of the BADST-2 that you wou	uld change	?			
5.	Is there anything you would add to the BADS	Γ-2?				
6	W hat other tools or resources would you like	to see?				
-	Trial office tools of resources would you like					

Please return this form to jon bull@sandwellchildrenstrust.org





The BADST-2 and the CRAST with the accompanying Short Brief Intervention Tools (SBIT's) for Alcohol, Drugs, and Wellbeing can be downloaded in pdf format at www.ourguideto.co.uk where you will also find the DECCA Awareness Club and further resources.

For alcohol and drugs information, news, and further resources go to: www.ourguideto.co.uk



Further training in the use of these tools and the Short Brief Intervention Tools is available to book on the website of the Sandwell Children's Safeguarding Partnership (SCSP)

https://training.sandwelllscb.org.uk/



If you live in Sandwell and are thinking about reducing the amount you are drinking, you can also access an app which can identify how much you are drinking and offers advice on how you can cut down.

The <u>Lower My Drinking app</u> can be downloaded through Google Play or iTunes, it is available in English and Polish.



As <u>GamCare partner</u> in the Midlands, Aquarius provide support, information and advice to anyone suffering with a gambling problem, as well as to family members and friends affected by someone else's gambling.

Sandwell Adult Social Care
Oldbury Council House
Freeth Street
PO BOX 2374
Oldbury
B69 3DE

www.sandwell.gov.uk